



## Academic eferences

Adamsen L, Quist M, Andersen C, et al. Effect of a multimodal high intensity exercise intervention in cancer patients undergoing chemotherapy: randomised controlled trial. *BMJ*. 2009; 339:b3410. [PMC free article] [PubMed] [Google Scholar]

Ainsworth BE, Haskell WL, Whitt MC, et al. Compendium of physical activities: an update of activity codes and MET intensities. *Med Sci Sports Exerc*. 2000;32(9 Suppl):S498–S504. [PubMed] [Google Scholar]

Akechi T, Kugaya A, Okamura H, Yamawaki S, Uchitomi Y. Fatigue and its associated factors in ambulatory cancer patients: a preliminary study. *J Pain Symptom Manage*. 1999;17(1):42–48. [PubMed] [Google Scholar]

Baguely, BJ, Bolam, KA, Wright OR, and Skinner TL. The Effect of Nutrition Therapy and Exercise on Cancer-Related Fatigue and Quality of Life in Men with Prostate Cancer: A Systematic Review. *Nutrients*. 2017; 9(9), 1003 <https://doi.org/10.3390/nu9091003> (PubMed) (Google Scholar)

Begg CB. Publication bias. In: Cooper H, Hedges LV, editors. *The Handbook of Research Synthesis*. New York: Russell Sage Foundation; 1994. pp. 399–409. [Google Scholar]

Borenstein M, Hedges LV, Higgins JPT, Rothstein HR. *Comprehensive Meta-analysis (Version 22027)* [Computer software] Englewood, NJ: Biostat; 2005. [Google Scholar]

Borenstein M, Hedges LV, Higgins JPT, Rothstein HR. *Introduction to Meta-analysis (Statistics in Practice)* Chichester, UK: John Wiley & Sons; 2009. [Google Scholar]

Bower JE, Lamkin DM. Inflammation and cancer-related fatigue: mechanisms, contributing factors, and treatment implications. *Brain Behav Immun*. 2013;30(Suppl): S48–S57. [PMC free article] [PubMed] [Google Scholar]

Cancer Research UK Worldwide cancer incidence statistics. 2014. Available from: <http://www.cancerresearchuk.org/cancer-info/cancerstats/world/incidence>

Cantarero-Villanueva I, Fernández-Lao C, Cuesta-Vargas AI, Del Moral-Avila R, Fernández-de-Las-Peñas C, Arroyo-Morales M. The effectiveness of a deep-water aquatic exercise program in cancer-related fatigue in breast cancer survivors: a randomized controlled trial. *Arch Phys Med Rehabil*. 2013;94(2):221–230. [PubMed] [Google Scholar]

Cella D, Davis K, Breitbart W, Curt G, Fatigue Coalition Cancer-related fatigue: prevalence of proposed diagnostic criteria in a United States sample of cancer survivors. *J Clin Oncol*. 2001;19(14):3385–3391. [PubMed] [Google Scholar]

Cohen J. *Statistical Power Analysis for the Behavioral Sciences*. 2nd ed. Hillsdale, NJ: Lawrence Erlbaum Associates; 1988. [Google Scholar]

Courneya KS, Friedenreich CM, Quinney HA, et al. A longitudinal study of exercise barriers in colorectal cancer survivors participating in a randomized controlled trial. *Ann Behav Med*. 2005;29(2):147–153. [PubMed] [Google Scholar]

Cramp F, Byron-Daniel J. Exercise for the management of cancer-related fatigue in adults. *Cochrane Database Syst Rev*. 2012;(11):CD006145. [PubMed] [Google Scholar]

Dodd MJ, Cho MH, Miaskowski C, et al. A randomized controlled trial of home-based exercise for cancer-related fatigue in women during and after chemotherapy with or without radiation therapy. *Cancer Nurs*. 2010;33(4):245–257. [PMC free article] [PubMed] [Google Scholar]

Donnelly CM, Blaney JM, Lowe-Strong A, et al. A randomised controlled trial testing the feasibility and efficacy of a physical activity behavioural change intervention in managing fatigue with gynaecological cancer survivors. *Gynecol Oncol*. 2011;122(3):618–624. [PubMed] [Google Scholar]

Drewnowski A, Evans WJ. Nutrition, physical activity, and quality of life in older adults: summary. *J Gerontol A Biol Sci Med Sci*. 2001;89–94. 56 Spec 2. [PubMed] [Google Scholar]

Fukuda K, Straus SE, Hickie I, Sharpe MC, Dobbins JG, Komaroff A. The chronic fatigue syndrome: a comprehensive approach to its definition and study. International Chronic Fatigue Syndrome Study Group. *Ann Intern Med*. 1994;121(12):953–959. [PubMed] [Google Scholar]

Hall AE, Boyes AW, Bowman J, Walsh RA, James EL, Girgis A. Young adult cancer survivors' psychosocial well-being: a cross-sectional study assessing quality of life, unmet needs, and health behaviors. *Support Care Cancer*. 2012;20(6):1333–1341. [PubMed] [Google Scholar]

Higgins JP, Altman DG, Gøtzsche PC, et al. Cochrane Bias Methods Group; Cochrane Statistical Methods Group The Cochrane Collaboration's tool for assessing risk of bias in randomised trials. *BMJ*. 2011; 343:d5928. [PMC free article] [PubMed] [Google Scholar]

Higgins JP, Thompson SG, Deeks JJ, Altman DG. Measuring inconsistency in meta-analyses. *BMJ*. 2003;327(7414):557–560. [PMC free article] [PubMed] [Google Scholar]

Hofman M, Morrow GR, Roscoe JA, et al. Cancer patients' expectations of experiencing treatment-related side effects: a University of Rochester Cancer Center–Community Clinical

Oncology Program study of 938 patients from community practices. *Cancer*. 2004;101(4):851–857. [PubMed] [Google Scholar]

Humpel N, Iverson DC. Depression and quality of life in cancer survivors: is there a relationship with physical activity? *Int J Behav Nutr Phys Act*. 2007; 4:65. [PMC free article] [PubMed] [Google Scholar]

Husebø AM, Karlsen B, Allan H, Søreide JA, Bru E. Factors perceived to influence exercise adherence in women with breast cancer participating in an exercise programme during adjuvant chemotherapy: a focus group study. *J Clin Nurs*. 2015;24(3–4):500–510. [PubMed] [Google Scholar]

Jones LW, Courneya KS, Vallance JK, et al. Understanding the determinants of exercise intentions in multiple myeloma cancer survivors: an application of the theory of planned behavior. *Cancer Nurs*. 2006;29(3):167–175. [PubMed] [Google Scholar]

Kellner R, Sheffield BF. The one-week prevalence of symptoms in neurotic patients and normals. *Am J Psychiatry*. 1973; 50:734–741. [PubMed] [Google Scholar]

Kempke S, Goossens L, Luyten P, Bekaert P, Van Houdenhove B, Van Wambeke P. Predictors of outcome in a multi-component treatment program for chronic fatigue syndrome. *J Affect Disord*. 2010;126(1–2):174–179. [PubMed] [Google Scholar]

Kessels E, Husson O, and van der Feltz-Cornelis, CM, The effect of exercise on cancer-related fatigue in cancer survivors: a systematic review and meta-analysis. *Neuropsychiatric Disease and Treatment*. 2018; 14: 479-494. (PubMed) (Google Scholar)

Kessels E, Husson O, van der Feltz-Cornelis CM. Review protocol. 2017. [Accessed January 10, 2018]. Available from: [http://www.crd.york.ac.uk/PROSPERO/display\\_record.asp?ID=CRD42013003670](http://www.crd.york.ac.uk/PROSPERO/display_record.asp?ID=CRD42013003670).

Larson-Meyer DE. A systematic review of the energy cost and metabolic intensity of yoga. *Med Sci Sports Exerc*. 2016;48(8):1558–1569. [PubMed] [Google Scholar]

Larun L, Brurberg KG, Odgaard-Jensen J, Price JR. Exercise therapy for chronic fatigue syndrome. *Cochrane Database Syst Rev*. 2015;(2):CD003200. [PubMed] [Google Scholar]

McArdle William D, Katch Frank I, Katch Victor L. *Essentials of Exercise Physiology*. Vol. 2006. Philadelphia: Lippincott Williams & Wilkins; p. 204. [Google Scholar]

Mendoza TR, Wang XS, Cleeland CS, et al. The rapid assessment of fatigue severity in cancer patients: use of the Brief Fatigue Inventory. *Cancer*. 1999;85(5):1186–1196. [PubMed] [Google Scholar]

Midtgaard J, Baadsgaard MT, Møller T, et al. Self-reported physical activity behaviour; exercise motivation and information among Danish adult cancer patients undergoing chemotherapy. *Eur J Oncol Nurs*. 2009;13(2):116–121. [PubMed] [Google Scholar]

Mock V, Frangakis C, Davidson NE, et al. Exercise manages fatigue during breast cancer treatment: a randomized controlled trial. *Psychooncology*. 2005;14(6):464–477. [PubMed] [Google Scholar]

Moher D, Liberati A, Tetzlaff J, Altman DG, PRISMA Group Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *J Clin Epidemiol*. 2009;62(10):1006–1012. [PubMed] [Google Scholar]

Mustian KM, Alfano CM, Heckler C, et al. Comparison of pharmaceutical, psychological, and exercise treatments for cancer-related fatigue: a meta-analysis. *JAMA Oncol*. 2017;3(7):961–968. [PMC free article] [PubMed] [Google Scholar]

Mustian KM, Peppone L, Darling TV, Palesh O, Heckler CE, Morrow GR. A 4-week home-based aerobic and resistance exercise program during radiation therapy: a pilot randomized clinical trial. *J Support Oncol*. 2009;7(5):158–167. [PMC free article] [PubMed] [Google Scholar]

NCCN Clinical Practice Guidelines in Oncology Cancer-related fatigue. 2017. (Version 2). Available from: [https://www.nccn.org/professionals/physician\\_gls/pdf/fatigue.pdf](https://www.nccn.org/professionals/physician_gls/pdf/fatigue.pdf).

Pachman DR, Barton DL, Swetz KM, Loprinzi CL. Troublesome symptoms in cancer survivors: fatigue, insomnia, neuropathy, and pain. *J Clin Oncol*. 2012;30(30):3687–3696. [PubMed] [Google Scholar]

Patrick DL, Ferketich SL, Frame PS, et al. National Institutes of Health State-of-the-Science Panel National Institutes of Health State-of-the-Science Conference Statement: symptom management in cancer: pain, depression, and fatigue, July 15–17, 2002. *J Natl Cancer Inst Monogr*. 2004;(32):9–16. [PubMed] [Google Scholar]

Peterson MD, Gordon PM. Resistance exercise for the ageing adult: clinical implications and prescription guidelines. *Am J Med*. 2011;124(3):194–198. [PubMed] [Google Scholar]

Piotrowicz E, Orzechowski P, Chrapowicka A, Piotrowicz R. How should we advise heart failure patients on exercise and what should we tell them? *Curr Heart Fail Rep*. 2014;11(3):274–280. [PubMed] [Google Scholar]

Plowman Sharon A, Smith Denise L. *Exercise Physiology for Health, Fitness, and Performance*. Philadelphia: Lippincott Williams & Wilkins; 2001. p. 61. [Google Scholar]

Rogers LQ, Vicari S, Trammell R, et al. Biobehavioral factors mediate exercise effects on fatigue in breast cancer survivors. *Med Sci Sports Exerc*. 2014;46(6):1077–1088. [PMC free article] [PubMed] [Google Scholar]

Seo Y, Oh H, Seo W. Causal relationships among factors associated with cancer-related fatigue. *Eur J Oncol Nurs*. 2010;14(5):380–386. [PubMed] [Google Scholar]

Servaes P, Gielissen MF, Verhagen S, Bleijenberg G. The course of severe fatigue in disease-free breast cancer patients: a longitudinal study. *Psychooncology*. 2007;16(9):787–795. [PubMed] [Google Scholar]

Servaes P, van der Werf S, Prins J, Verhagen S, Bleijenberg G. Fatigue in disease-free cancer patients compared with fatigue in patients with chronic fatigue syndrome. *Support Care Cancer*. 2001;9(1):11–17. [PubMed] [Google Scholar]

Sharpe M, Goldsmith KA, Johnson AL, Chalder T, Walker J, White PD. Rehabilitative treatments for chronic fatigue syndrome: long-term follow-up from the PACE trial. *Lancet Psychiatry*. 2015;2(12):1067–1074. [PubMed] [Google Scholar]

Son CG. Review of the prevalence of chronic fatigue worldwide. *J Korean Orient Med*. 2012;33(2):25–33. [Google Scholar]

US National Coalition for Cancer Survivorship Defining cancer survivorship. Available from: <http://www.canceradvocacy.org/news/defining-cancer-survivorship>

Van Houdenhove B, Bruyninckx K, Luyten P. In search of new balance. Can high “action-proneness” in patients with chronic fatigue syndrome be changed by a multidisciplinary group treatment? *J Psychosom Res*. 2006;60(6):623–625. [PubMed] [Google Scholar]

van Houdenhove B, Heijnen CJ. [Chronic fatigue syndrome: a psychoneuroimmunological perspective] *Tijdschr Psychiatr*. 2009;51(8):603–610. Dutch. [PubMed] [Google Scholar]

Van Houdenhove B, Luyten P. Chronic fatigue syndrome reflects loss of adaptability. *J Intern Med*. 2010;268(3):249–251. [PubMed] [Google Scholar]

Van Houdenhove B, Van Den Eede F, Luyten P. Does hypothalamic-pituitary-adrenal axis hypofunction in chronic fatigue syndrome reflect a ‘crash’ in the stress system? *Med Hypotheses*. 2009;72(6):701–705. [PubMed] [Google Scholar]

Wang XS, Zhao F, Fisch MJ, et al. Prevalence and characteristics of moderate to severe fatigue: a multicenter study in cancer patients and survivors. *Cancer*. 2014;120(3):425–432. [PMC free article] [PubMed] [Google Scholar]

Wang YJ, Boehmke M, Wu YW, Dickerson SS, Fisher N. Effects of a 6-week walking program on Taiwanese women newly diagnosed with early-stage breast cancer. *Cancer Nurs*. 2011;34(2):E1–E13. [PubMed] [Google Scholar]

Wessely S, Powell R. Fatigue syndromes: a comparison of chronic “postviral” fatigue with neuromuscular and affective disorders. *J Neurol Neurosurg Psychiatry*. 1989;52(8):940–948. [PMC free article] [PubMed] [Google Scholar]

Williams TE, Chalder T, Sharpe M, White PD. Heterogeneity in chronic fatigue syndrome – empirically defined subgroups from the PACE trial. *Psychol Med*. 2017:1–12. [PubMed] [Google Scholar]

Williams TE, Chalder T, Sharpe M, White PD. Heterogeneity in chronic fatigue syndrome – empirically defined subgroups from the PACE trial. *Psychol Med*. 2017;47(8):1454–1465. [PubMed] [Google Scholar]

Windsor PM, Nicol KF, Potter J. A randomized, controlled trial of aerobic exercise for treatment-related fatigue in men receiving radical external beam radiotherapy for localized prostate carcinoma. *Cancer*. 2004;101(3):550–557. [PubMed] [Google Scholar]

Wu HS, McSweeney M. Cancer-related fatigue: “It’s so much more than just being tired” *Eur J Oncol Nurs*. 2007;11(2):117–125. [PubMed] [Google Scholar]

Yeo TP, Burrell SA, Sauter PK, et al. A progressive postresection walking program significantly improves fatigue and health-related quality of life in pancreas and periampullary cancer patients. *J Am Coll Surg*. 2012;214(4):463–475. discussion 475–477. [PubMed] [Google Scholar]

Yuen HK, Sword D. Home-based exercise to alleviate fatigue and improve functional capacity among breast cancer survivors. *J Allied Health*. 2007;36(4):e257–e275. [PubMed] [Google Scholar]